



Brief paper

A result on second order nonlinear operators arising in high-speed networking applications[☆]

Martin Corless^{a,*}, Chris King^b, Robert Shorten^c

^a School of Aeronautics & Astronautics, Purdue University, USA

^b Department of Mathematics, Northeastern University, USA

^c The Hamilton Institute, NUI Maynooth, Ireland

ARTICLE INFO

Article history:

Received 5 April 2008

Received in revised form

11 July 2008

Accepted 8 December 2008

Available online 6 March 2009

Keywords:

Communication networks

Congestion control

TCP

Nonlinear systems

ABSTRACT

We consider the behaviour of a class of nonlinear difference equations that arise in networks employing congestion control protocols. For a class of such systems we show that trajectories, experiencing the same set of congestion notifications, but starting at different initial conditions, converge exponentially to each other.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Recent years have witnessed a huge interest in both the analysis and design of congestion control protocols for deployment in the internet (Low, Paganini, & Doyle, 2002). In particular, protocols based on the *Additive-Increase Multiplicative-Decrease* (AIMD) paradigm, of which TCP (Jacobson, 1988) is one, have been the subject of much scrutiny. The study of such protocols is interesting for a number of reasons. Clearly, given the importance of the internet, the study of network congestion control is of immense practical value. Secondly, network congestion control brings together many concepts and ideas from diverse areas of Applied Mathematics. These include (to name but a few): Probability Theory (Barnsley, Demko, Elton, & Geronimo, 1988; Stenflo, 2002); the theory of Stochastic Matrices (Shorten, Wirth, & Leith, 2006); Positive Systems and Hybrid Systems (Baccelli & Hong, 2002). Many practical problems that arise in the study of congestion control therefore motivate mathematical questions that are not only of interest in the context of the application, but also merit study in a more general and abstract setting. Our

problem is motivated by such generalisations of the *Transmission Control Protocol* (TCP). Standard TCP is based on the AIMD algorithm of Chiu and Jain. In their original paper (Chiu & Jain, 1989), Chiu and Jain consider a system in which n users compete for a resource. The users' actions consist of (continuously) probing the availability of the resource by submitting requests for its use. A key assumption in the model is that the users do not communicate directly and that the only information they have about the availability of the resource is when the collective utilization of the resource reaches some capacity constraint. At such time instances, referred to as *congestion events*, some or all users are informed through feedback and respond instantly by decentralized down-scaling of their individual utilization rates. Given this basic setting, the problem is then to develop an algorithm that produces probing strategies for the users so that each user will infer its "fair" share of the shared resource in a decentralized manner, while at the same time preventing congestion collapse (Low et al., 2002).

The AIMD algorithm of Chiu and Jain describes probing strategies that evolve in cycles, each cycle having two phases (Rothblum & Shorten, 2007). The first phase is instantaneous. It occurs when capacity is reached, a subset of users are notified, and notified users respond by down-scaling their utilization rates (abruptly) by a multiplicative factor. During the second phase of a cycle, each user increases its utilization rate linearly until congestion is reached again, at which point the first phase of the next cycle is entered. Let $w_i(k) \geq 0$ be the i th user's share of the resource at the beginning of the k th cycle. Then,

$$w_i(k+1) = b_i(k)w_i(k) + \alpha_i T(k), \quad (1)$$

[☆] This paper was not presented at any IFAC meeting. This paper was recommended for publication in revised form by Associate Editor Masayuki Fujita under the direction of Editor Ian R. Petersen.

* Corresponding author. Tel.: +1 765 494 7411; fax: +1 765 494 0307.

E-mail addresses: corless@purdue.edu (M. Corless), c.king@neu.edu (C. King), Robert.Shorten@nuim.ie (R. Shorten).

where $T(k)$ is the time taken by the k th cycle; we call this the *inter-congestion time*. The positive scalar α_i is the rate of increase in the second phase. The scalar $b_i(k)$ depends on whether or not the i th user is informed of congestion at the k th event; if it is informed then $b_i(k)$ equals some constant β_i (call it the drop parameter) which satisfies $0 \leq \beta_i < 1$; if not, $b_i(k) = 1$. If C is the total capacity of the resource available then, at congestion,

$$w_1(k) + \dots + w_n(k) = w_1(k+1) + \dots + w_n(k+1) = C. \quad (2)$$

The inter-congestion time T is determined by (1) and (2).

1.1. Previous results and contribution

A primary objective of network congestion control is to prevent network congestion collapse, or more generally to maintain a reasonable throughput of useful information for users (as opposed to resending the same information that is continually being dropped). A second, equally important, objective is to ensure a fair and equitable allocation of network resources (bandwidth) in a distributed manner without communication between network users.

For this dynamical system it is essential to settle questions such as the existence and uniqueness of a trajectory which is independent of initial conditions, and the rate at which the system approaches this steady state trajectory. These convergence issues have been studied extensively for the standard AIMD-TCP model described above, and a complete picture for this model is presented in Shorten, King, and Wirth (2007). Even under very general assumptions, questions relating to the existence and uniqueness of network equilibria have been settled and are reported in Shorten et al. (2007).

Our objective in the present paper is to extend these results to generalisations of standard TCP. Such generalisations have been motivated by the view that standard TCP is deficient in most modern network scenarios (Floyd, 2003; Jin, Wei, & Low, 2003; Kelly, 2002; Xu, Harfoush, & Rhee, 2004). Recently nonlinear AIMD variants (NAIMD) have been proposed for deployment in networks. Remarkably, despite the fact that nonlinear TCPs are widely deployed in the internet (cubic is the default in Linux), few mathematical results exist that characterise their behaviour (one way or the other).

Generally speaking, NAIMD protocols differ from the standard (linear) AIMD in the second (AI) phase, whereby the linear increase in the utilization rate of each user i is replaced by a nonlinear function of time that we denote by a_i . Thus a user that is informed of congestion at the k th congestion event evolves as follows:

$$w_i(k+1) = \beta_i w_i(k) + a_i(T(k)). \quad (3)$$

However for a user that is not informed of congestion, the future evolution of its utilization rate depends in addition on how long it has been growing in the second phase. Specifically let $t_i(k)$ be the time elapsed since user i was last informed of a congestion event. Then the evolution equation for its utilization rate is

$$w_i(k+1) = w_i(k) + a_i(t_i(k) + T(k)) - a_i(t_i(k)). \quad (4)$$

Thus a complete description of the state at the k th congestion event must include the times t_i as well as the utilization rates w_i . So the state X of the system at a congestion event is a vector in \mathbb{R}^{2n} , namely

$$X = (w, t) \quad \text{where } w = (w_1, \dots, w_n) \text{ and } t = (t_1, \dots, t_n).$$

Also Eqs. (3) and (4) must be supplemented by the evolution equations for t_i . If user i is informed of congestion then

$$t_i(k+1) = T(k) \quad (5)$$

while if user i is not informed of congestion then

$$t_i(k+1) = t_i(k) + T(k). \quad (6)$$

Together (3)–(6) determine the evolution of the state vector $X(k) \mapsto X(k+1)$ between successive congestion events.

Initial attempts to study general nonlinear AIMD models are presented in Rothblum and Shorten (2007) and King, Shorten, Wirth, and Akar (2008); albeit for the case of synchronised communication networks. Motivated by the desire to ultimately extend this work to a stochastic setting mimicking that in Shorten et al. (2007), we present here in this present paper an initial analysis of nonlinear AIMD protocols which we call *eventually linear*. These models are characterised by growth functions a_i which become (affine) linear after some time T_{m_i} . Specifically, such a growth function a_i satisfies

$$a_i(\tau') = a_i(T_{m_i}) + \alpha_{H_i}(\tau' - T_{m_i}) \quad \text{for } \tau' \geq T_{m_i} \quad (7)$$

where α_{H_i} is constant, and where T_{m_i} is the ‘elbow’ in the function. So after time T_{m_i} in the second phase the model reverts to standard linear TCP with utilization rate growing at the rate α_{H_i} . For the synchronised case it is known that all solutions converge to a unique fixed point at a geometrical rate (Rothblum & Shorten, 2007). Here we begin the analysis of the unsynchronised case of the eventually linear model, by addressing the question of how the solution depends on the initial value $w(0)$ for the user shares.

2. Eventually linear congestion control protocols

The main motivation for moving away from linear AIMD is to realise protocols whose properties scale with increasing network bandwidths. This requirement suggests developing algorithms whose probing phase is nonlinear or piecewise linear in nature, where each linear phase is designed with a typical network (bandwidth) scenario in mind. In this latter setting, it is entirely reasonable to assume that it only rarely occurs that network flows do not achieve the phase of the probing action that is suitable for current network conditions. From a mathematical perspective, this is a very appealing assumption. In essence, this means that while eventually linear protocols are nonlinear and thus include all the complexities of NAIMD, including long-term memory effects, this latter assumption means that the nonlinear features (which dominate only when the inter-congestion times are short) can be treated fully without approximation in this regime. In particular, if all inter-congestion times are greater than a certain minimum value then the growth functions of every flow move past the ‘elbow’ before the next congestion event occurs. This latter fact means that the equations describing the difference dynamics are the same as in standard TCP and all our previous results apply. In what follows we explore this intuition in a more formal setting. We begin by defining the notion of a *regular* congestion event.

Definition 2.1. A congestion event k is *regular* if $T(k) \geq T_{m_i}$ for every user i .

Thus a regular congestion event occurs whenever the subsequent inter-congestion time exceeds all the times T_{m_i} , meaning that every flow enters its linear phase before the next congestion event. The essential feature of regular events is that they ensure that all memory of the past is lost and the network dynamics are in effect linear. To explore the consequences of regular events we will compare two trajectories, both experiencing the same selection of dropped flows at each congestion event, but evolving from two different initial states. Our principal assumption is that the congestion events are all regular for both trajectories. We denote the set of states of the first trajectory at successive congestion

events by $\{w(0), w(1), \dots, w(k)\}$, where $w(0)$ is the initial condition, and the set of states associated with the second trajectory by $\{v(0), v(1), \dots, v(k)\}$. Denote further the difference vectors by $E(k) = w(k) - v(k)$. Then, the assumption that all events are regular implies that the difference vectors evolve linearly between the $t_i(k)$, that is,

$$\begin{aligned} E_i(k+1) &= w_i(k+1) - v_i(k+1) \\ &= b_i(k)(w_i(k) - v_i(k)) + \alpha_{H_i}(T(k) - S(k)) \end{aligned} \quad (8)$$

where $b_i(k) = \beta$ or $b_i(k) = 1$ depending on whether or not this flow experiences a drop at the event, and where $T(k), S(k)$ are the inter-congestion times for the w, v flows respectively. Since both $E(k)$ and $E(k+1)$ are perpendicular to the vector $e = [1, 1, \dots, 1]^T$ this set of equations determines the value of $T(k) - S(k)$, and leads to the linear relation

$$E(k+1) = A(k)E(k), \quad A(k) \in \mathcal{A} \quad (9)$$

where \mathcal{A} is precisely a set of matrices obtained from the dynamics of standard unsynchronised TCP. In particular, each $A(k)$ is entrywise non-negative and column stochastic and hence non-expansive in the one-norm; thus $\|E(k+1)\| \leq \|E(k)\|$ for all k . Here and throughout the paper $\|\cdot\|$ denotes the one-norm of a vector, that is, the sum of the absolute values of its components.

Definition 2.2. A drop sequence is *typical* if there is an integer N such that every flow experiences at least one drop during N successive congestion events.

We then have the following theorem.

Theorem 2.1. *Let $w(0), v(0)$ be any initial conditions, and consider a typical drop sequence for which both flow sequences $\{w(k)\}$ and $\{v(k)\}$ experience only regular congestion events. Then $\|E(k)\| = \|w(k) - v(k)\|$ converges to zero at an exponential rate as $k \rightarrow \infty$, and the rate is uniform in $w(0), v(0)$.*

Proof. It follows from (9) that

$$E(k) = A(k)A(k-1) \dots A(0)E(0). \quad (10)$$

Since the drop sequence is typical there is an integer N such that every flow experiences a drop at least once in every N events. Let $k = \kappa N + l$ where $0 \leq l \leq N-1$ is the remainder, and define for $j = 0, \dots, \kappa-1$

$$B(j) = A(jN + N - 1)A(jN + N - 2) \dots A(jN). \quad (11)$$

Then

$$E(k) = A(\kappa N + l) \dots A(\kappa N)B(\kappa - 1) \dots B(1)B(0)E(0).$$

Each matrix $A(k)$ is non-expansive on the hyperplane in \mathbb{R}^n orthogonal to $[1, 1, \dots, 1]^T$, that is $\|A(k)E\| \leq \|E\|$, for all E in this hyperplane. Each matrix $B(j)$ is column stochastic and entrywise positive. This follows from the assumption that the drop sequence is regular, and from the fact that the columns of $A(k)$ corresponding to its dropped flows are entrywise positive. It therefore follows that each matrix $B(j)$ is a contraction on the hyperplane in \mathbb{R}^n orthogonal to $[1, 1, \dots, 1]^T$. Since the set \mathcal{A} is finite (it has at most $2^n - 1$ elements), it follows that there is there is a finite number of different $B(j)$ matrices (at most $(2^n - 1)^N$). Hence there is $r < 1$ such that for all j and for all E orthogonal to $[1, 1, \dots, 1]^T$,

$$\|B(j)E\| \leq r \|E\|. \quad (12)$$

It follows that

$$\|E(k)\| \leq r^\kappa \|E(0)\| \leq r^{k/N-1} \|E(0)\|. \quad (13)$$

which yields exponential convergence at rate $r^{1/N}$, uniform in $w(0), v(0)$. \square

Before proceeding to the statement of our main result, we present the following lemma (which is proven in the [Appendix](#)).

Lemma 2.1. *If $t_i(k^*) \geq T_{m_i}$ for every user i which is not dropped at some stage k^* then, $w(k)$ is independent of $t(k^*)$ for all $k \geq k^*$.*

As we shall see in the proof of our main result, this lemma is central to all that follows and with its help we shall prove that in spite of nonlinear and long-term memory effects, trajectories of two-flow systems starting from different initial conditions but experiencing the same drop sequence converge to each other exponentially, and that this is true for every possible drop sequence.

3. A convergence result for two users

We now present the main mathematical results of this paper. Our main result says that for a two-user network, trajectories starting from different initial conditions, but experiencing the same set of congestion events, converge exponentially to each other.

Setting: In what follows $w = (w_1, w_2)$ and $t = (t_1, t_2)$, and the evolution of the our dynamic system is described by the four difference equations

$$\begin{aligned} w_i(k+1) &= b_i w_i + a_i(d_i t_i + T) - a_i(d_i t_i) \\ t_i(k+1) &= d_i t_i + T \end{aligned} \quad i = 1, 2 \quad (14)$$

where, for simplicity of presentation, we sometimes omit the dependence on k of variables on the right-hand side of a difference equation. The scalars $b_i(k)$ and $d_i(k)$ are defined as follows: if the i th user is dropped at event k then, $(b_i(k), d_i(k)) = (\beta_i, 0)$; if the i th user is not dropped at stage k then, $(b_i(k), d_i(k)) = (1, 1)$. As usual, the time between congestion events T , is determined by the congestion condition that

$$w_1(k) + w_2(k) = w_1(k+1) + w_2(k+1) = C$$

and system (14) evolves on the extended state space

$$\mathcal{S}^{\text{ext}} = \{(w, t) : w \in \mathcal{S}, t_1, t_2 \geq 0\} \quad (15)$$

where

$$\mathcal{S} = \{w \in \mathbb{R}^2 : w_1, w_2 \geq 0 \text{ and } w_1 + w_2 \leq C\} \quad (16)$$

is the rate space.

We assume that both users have growth rate functions which are eventually affine as described in (7). We also assume that each a_i is *increasing and convex* on $[0, \infty)$, $a_i(0) = 0$ and there exists $\alpha_{L_i} > 0$ such that $a_i(\tau_2) - a_i(\tau_1) \geq \alpha_{L_i}(\tau_2 - \tau_1)$ for all $\tau_2 \geq \tau_1$.

Statement of the main result: For the special case of two users we show that for a given drop sequence, all trajectories converge exponentially to a unique trajectory.

The solution of system (14) is determined by the initial state $(w(0), t(0))$ and by the sequence of drops at the congestion events. Our main result is that when the system capacity C is sufficiently large, the long-term behaviour of w is determined solely by the sequence of drops and is independent of the initial state.

Consider first the cases in which one of the users is never dropped. If the first user is never dropped, one can show that, regardless of initial conditions, $\lim_{k \rightarrow \infty} w(k) = (1, 0)$. For the second user, the consequence is $\lim_{k \rightarrow \infty} w(k) = (0, 1)$. We call the drop sequences corresponding to these cases trivial.

Our main result requires that C, T_{m_1}, T_{m_2} satisfy the following condition.

$$C \geq \max\{C_0, C_1, C_2\} \quad (17)$$

where

$$C_0 = w_1 + w_2 \quad (18)$$

$$C_1 = \frac{(\alpha_{H_1} + \alpha_{H_2})w_1 + \alpha_{H_1}a_2(T_m) - \alpha_{H_2}a_1(T_m)}{(1 - \beta_2)\alpha_{H_1}} \quad (19)$$

$$C_2 = \frac{(\alpha_{H_1} + \alpha_{H_2})w_2 + \alpha_{H_2}a_1(T_m) - \alpha_{H_1}a_2(T_m)}{(1 - \beta_1)\alpha_{H_2}} \quad (20)$$

with

$$w_1 = \frac{a_1(T_m) + \alpha_{H_2}T_m}{1 - \beta_1}, \quad w_2 = \frac{a_2(T_m) + \alpha_{H_1}T_m}{1 - \beta_2} \quad (21)$$

and

$$T_m = \max\{T_{m_1}, T_{m_2}\}. \quad (22)$$

Note that C_0, C_1, C_2 go to zero as T_{m_1}, T_{m_2} go to zero. Hence, for a given system capacity C , condition (17) can be satisfied by choosing T_{m_1} and T_{m_2} sufficiently small.

We now present the main result of the paper.

Theorem 3.1. *When condition (17) is satisfied, there is a constant $r < 1$ such that the following holds for every non-trivial drop sequence. There are constants $c > 0$ and k^* such that if (w, t) and (v, s) are any two solutions of (14) then,*

$$\|w(k) - v(k)\| \leq cr^{k/2}(\|w(0) - v(0)\| + \|t(0) - s(0)\|)$$

for all $k \geq k^*$.

Theorem 3.1 shows that rate vector histories corresponding to different initial conditions exponentially converge onto a trajectory which is determined solely by the sequence of dropped users at the congestion events. Hence the long-run behaviour of the rate vector does not depend on the initial conditions.

Comment: Our main theorem does not exclude the possibility that the long-run behaviour could depend on how the dropped users are selected at the congestion events, since different sequences of drops will lead to different trajectories.

4. Proof of the main result

We now present the proof of our main result. An essential component of our proof is the existence of a reduced order dynamic system that captures the convergence properties of the original system.

4.1. Preamble to proof: A reduced order system

At a congestion event, there are only three possibilities: only user one is dropped, only user two is dropped or both users are dropped. We use the integer q_k to indicate the set of users dropped at an event k . We let $q_k = 1, 2$, or 3 corresponding to the three possibilities as listed in the above order. With $X = (w, t)$, the evolution of system (14) can then be described by

$$X(k+1) = \Psi_{q_k}(X(k)) \quad (23)$$

where $q_k = 1, 2$ or 3 and the maps Ψ_1, Ψ_2, Ψ_3 are determined from (14) by choosing appropriate values of b_i and d_i . For example, Ψ_1 is obtained by letting $(b_1, d_1) = (\beta_1, 0)$ and $(b_2, d_2) = (1, 1)$ in (14). We also introduce the following two-step maps

$$\Psi_4 = \Psi_2 \circ \Psi_1, \quad \Psi_5 = \Psi_1 \circ \Psi_2. \quad (24)$$

Consider the following five subsets of the extended state space:

$$\mathcal{M}_1 = \{X \in \mathcal{S}^{\text{ext}} : t_2 \geq T_{m_2}\} \quad (25)$$

$$\mathcal{M}_2 = \{X \in \mathcal{S}^{\text{ext}} : t_1 \geq T_{m_1}\} \quad (26)$$

$$\mathcal{M}_3 = \mathcal{S}^{\text{ext}}, \quad \mathcal{M}_4 = \mathcal{M}_1, \quad \mathcal{M}_5 = \mathcal{M}_2. \quad (27)$$

For $p = 1, 2, 3$, the condition that $X \in \mathcal{M}_p$ is equivalent to the requirement that $t_i \geq T_{m_i}$ if the i th user is not dropped for the map Ψ_p . For $p = 4$ or 5 , $X \in \mathcal{M}_p$ means that $t_i \geq T_{m_i}$ if the i th user is not dropped for the first stage of the map Ψ_p .

As we have stated our objective here is to show that our original system is equivalent in some sense to a reduced order system. We establish this equivalence using Lemmas 4.1–4.4 and Corollaries 4.1, 4.2. When not explicitly given, proofs are presented in the Appendix. The first of these results is an immediate consequence of Lemma 2.1.

Lemma 4.1. *For $p = 1, \dots, 5$, $\Psi_p(X)$ is independent of t when $X \in \mathcal{M}_p$.*

Now let P be the projection of \mathcal{S}^{ext} onto \mathcal{S} defined by

$$P(w, t) = w. \quad (28)$$

For $p = 1, \dots, 5$, it follows from the last result that if we define the map Φ_p on the rate space \mathcal{S} by

$$\Phi_p(w) = P\Psi(w, T_m, T_m) \quad (29)$$

where T_m is defined in (22) then,

$$P\Psi_p(X) = \Phi_p(PX) \quad \text{for all } X \in \mathcal{M}_p. \quad (30)$$

Recalling (21) we have the following result and corollary.

Lemma 4.2. *If a user i^* is dropped at an event k where $w_{i^*}(k) \geq \underline{w}_{i^*}$ then, $T(k) \geq T_m$ and $t_i(k+1) \geq T_m$ for $i = 1, 2$.*

In particular, the following is an immediate consequence of the previous discussion.

Corollary 4.1. *If both users are dropped at an event k then, $T(k) \geq T_m$ and $t_i(k+1) \geq T_m$ for $i = 1, 2$.*

Proof. Assumption (17) on C tells us that $C \geq C_0 = w_1 + w_2$. Since $w_1 + w_2 = C$ we must have $w_1 \geq \underline{w}_1$ or $w_2 \geq \underline{w}_2$. Since both users are dropped, it now follows from Lemma 4.2 that $T(k) \geq T_m$ and $t_i(k+1) \geq T_m$ for $i = 1, 2$. \square

Furthermore, the following lemma and corollary also follow.

Lemma 4.3. *If user i^* is not dropped at an event k then, $w_{i^*}(k+1) \geq \underline{w}_{i^*}$.*

Corollary 4.2. *If for some i^* , user i^* is not dropped at an event k but is dropped at event $k+1$ then, $T(k+1) \geq T_m$ and $t_i(k+2) \geq T_m$ for $i = 1, 2$.*

Proof. Since user i^* is not dropped at event k , Lemma 4.3 tells us that $w_{i^*}(k+1) \geq \underline{w}_{i^*}$. It now follows from Lemma 4.2 that $T(k+1) \geq T_m$ and $t_i(k+2) \geq T_m$ for $i = 1, 2$. \square

The following result motivates the introduction of Ψ_4 and Ψ_5 ; it follows from Corollaries 4.1 and 4.2. In this result

$$\mathcal{M} = \{X \in \mathcal{S}^{\text{ext}} : t_1, t_2 \geq T_m\}. \quad (31)$$

Note that $\mathcal{M} \subset \mathcal{M}_p$ for all p .

Lemma 4.4. *If $p = 1, \dots, 5$ then, \mathcal{M}_p is invariant for Ψ_p , that is, $\Psi_p(X) \in \mathcal{M}_p$ for $X \in \mathcal{M}_p$. If $p = 3, 4, 5$ then, $\Psi_p(X) \in \mathcal{M}$ for all $X \in \mathcal{S}^{\text{ext}}$.*

We now have the main result of this section.

Lemma 4.5 (Reduction Lemma). Suppose (w, t) is any solution of (14) and $t_i(k^*) \geq T_m$ for $i = 1, 2$ at some event k^* . Then for any event $k^f \geq k^*$, there is a subsequence $\{k_0, k_1, \dots, k_l\}$ of $\{k^*, k^* + 1, \dots, k^f\}$ with $k_0 = k^*$, $k_l = k^f$, $l \geq (k^f - k^*)/2$ and a sequence $\{p_0, p_1, \dots, p_{l-1}\}$ such that

$$w(k_{j+1}) = \Phi_{p_j}(w(k_j)) \quad (32)$$

for $j = 0, \dots, l-1$.

Remark 4.1. The above lemma tells us that if $t_1(k^*), t_2(k^*) \geq T_m$ at some event k^* then, the resulting behaviour of the rate vector w for the original full order system (14) is completely determined by the reduced order system (32).

4.2. Contractions and the proof of the main result

We now proceed to prove the main result of the paper. As before, proofs not explicitly given, are presented in the Appendix. Before we proceed let us now define the contraction parameter

$$r = \max\{r_1, r_2, r_4, r_5\}, \quad (33)$$

where

$$\begin{aligned} r_1 &= \frac{\alpha_{H_1} + \alpha_{L_2}\beta_1}{\alpha_{H_1} + \alpha_{L_2}}, & r_2 &= \frac{\alpha_{H_2} + \alpha_{L_1}\beta_2}{\alpha_{H_2} + \alpha_{L_1}} \\ r_4 &= \frac{\alpha_{L_1}(\alpha_{H_1}\beta_2 + \alpha_{H_2}\beta_1) + \alpha_{H_2}(\alpha_{H_1}(1 + \beta_1\beta_2 - \beta_1) + \alpha_{H_2}\beta_1)}{(\alpha_{L_1} + \alpha_{H_2})(\alpha_{H_1} + \alpha_{H_2})} \\ r_4 &= \frac{\alpha_{L_2}(\alpha_{H_2}\beta_1 + \alpha_{H_1}\beta_2) + \alpha_{H_1}(\alpha_{H_2}(1 + \beta_1\beta_2 - \beta_2) + \alpha_{H_1}\beta_2)}{(\alpha_{L_2} + \alpha_{H_1})(\alpha_{H_1} + \alpha_{H_2})}. \end{aligned}$$

Careful examination of the above expressions reveals that $r < 1$.

Our first result in this section states that Ψ_1, Ψ_2, Ψ_3 are globally Lipschitz and Φ_1, Φ_2, Φ_3 are contractive.

Lemma 4.6. (i) There is a constant c_1 such that, for all $X, Y \in \mathcal{S}^{\text{ext}}$,

$$\|\Psi_q(X) - \Psi_q(Y)\| \leq c_1 \|X - Y\| \quad \text{for } q = 1, 2, 3. \quad (34)$$

(ii) For all $w, v \in \mathcal{S}$,

$$\|\Phi_q(w) - \Phi_q(v)\| \leq r \|w - v\| \quad \text{for } q = 1, 2, 3, \quad (35)$$

where $r < 1$ is given by (33).

The next result states that the maps Φ_4 and Φ_5 are contractive.

Lemma 4.7. For all $w, v \in \mathcal{S}$,

$$\|\Phi_p(w) - \Phi_p(v)\| \leq r \|w - v\| \quad \text{for } p = 4, 5, \quad (36)$$

where $r < 1$ is given by (33).

Now, armed with these preliminary results we give the proof of the main result of the paper.

Proof of Theorem 3.1. Let $X = (w, t)$ and $Y = (v, s)$ be any two solutions of (14) corresponding to some non-trivial drop sequence $\{q_1, q_2, \dots\}$. Then there exists an event k^* such that either $q_{k^*-1} = 3$ or $\{q_{k^*-2}, q_{k^*-1}\}$ equals $\{1, 2\}$ or $\{2, 1\}$. It follows from Corollaries 4.1 and 4.2 that for any of the above occurrences,

$$t_i(k^*), s_i(k^*) \geq T_m$$

for $i = 1, 2$.

Consider any event $k = k^f > k^*$. It now follows from the Reduction Lemma (Lemma 4.5) that there is a subsequence $\{k_0, k_1, \dots, k_l\}$ of $\{k^*, k^* + 1, \dots, k^f\}$ with $k_0 = k^*$, $k_l = k^f$, $l \geq (k^f - k^*)/2$ and a sequence $\{p_0, \dots, p_{l-1}\}$ such that

$$w(k_{j+1}) = \Phi_{p_j}(w(k_j)) \quad \text{and} \quad v(k_{j+1}) = \Phi_{p_j}(v(k_j)) \quad (37)$$

for $j = 0, \dots, l-1$. Recalling the contraction results in Lemmas 4.6 and 4.7 we now obtain that

$$\|w(k_{j+1}) - v(k_{j+1})\| \leq r \|w(k_j) - v(k_j)\| \quad (38)$$

for $j = 0, \dots, l-1$. It now follows from (38), $k^f = k_l$, $k_0 = k^*$ and $l \geq (k^f - k^*)/2$ that

$$\begin{aligned} \|w(k^f) - v(k^f)\| &= \|w(k_l) - v(k_l)\| \leq r^l \|w(k_0) - v(k_0)\| \\ &\leq r^{(k^f - k^*)/2} \|w(k^*) - v(k^*)\|. \end{aligned}$$

Hence,

$$\|w(k^f) - v(k^f)\| \leq r^{(k^f - k^*)/2} \|w(k^*) - v(k^*)\|. \quad (39)$$

From Lemma 4.6 there exists a constant c_1 independent of $w(0), v(0), t(0), s(0)$ such that (34) holds. Since $X(k+1) = \Psi_{q_k}(X(k))$ and $Y(k+1) = \Psi_{q_k}(Y(k))$ where $q_k = 1, 2$ or 3 , we have

$$\|X(k+1) - Y(k+1)\| \leq c_1 \|X(k) - Y(k)\|$$

for all k . This implies that

$$\begin{aligned} \|w(k^*) - v(k^*)\| &\leq \|X(k^*) - Y(k^*)\| \leq c_1^{k^*} \|X(0) - Y(0)\| \\ &= c_1^{k^*} (\|w(0) - v(0)\| + \|t(0) - s(0)\|). \end{aligned}$$

This, combined with (39) yields

$$\|w(k^f) - v(k^f)\| \leq c r^{k^f/2} (\|w(0) - v(0)\| + \|t(0) - s(0)\|)$$

where $c = (c_1/r^{1/2})^{k^*}$. \square

5. Example

As we have already mentioned, our ultimate objective is to demonstrate the existence of a unique stationary measure for a wide class of nonlinear congestion control protocols. Roughly speaking, the existence of such a measure means that the long-term average behaviour of the network does not depend on initial conditions. In situations, where simulation is used as the principal tool for analysis, knowledge that the average behaviour of the network is independent of starting condition is a fundamental requirement. In what follows we give an example to illustrate our basic result.

Our basic setup is as follows. Two identical NAIMD flows compete for ten units of capacity. Upon notification of congestion a flow reduces its rate to half the value it had achieved before congestion. After congestion notified flows probe for available bandwidth with piecewise linear growth rate

$$a_i(\tau') = \begin{cases} \alpha_L \tau' & \text{if } 0 \leq \tau' \leq 1 \\ \alpha_L + \alpha_H(\tau' - 1) & \text{if } 1 \leq \tau' \end{cases}$$

where $\alpha_L = 1$ and $\alpha_H = 5$. In these examples, we abuse notation and let $w(\tau) = (w_1(\tau), w_2(\tau))$ be the flow rate vector at time $\tau \in [0, \infty)$.

Example 1 (A Random Drop Sequence). Here we generate a sequence of fifty congestion notifications using the random number generator in Matlab. We follow the evolution of two trajectories, each experiencing these identical congestion notifications, but starting from different initial conditions. Figs. 1 and 2 depict the evolution of $w(\tau)$ from $w(0) = (2, 8)$ and $w(0) = (5, 5)$, respectively; in both cases $t(0) = (0, 0)$. Finally, we show in Fig. 3 the distance between these trajectories as measured by the 1-norm as a function of congestion event.




Fig. 1. Example: Evolution of $w(\tau)$ from $w(0) = (2, 8)$.




Fig. 2. Example 1: Evolution of $w(\tau)$ from $w(0) = (5, 5)$.




Fig. 3. Example 1: Distance between flow trajectories. Note that the 1-norm does not decrease monotonically; but the convergence is exponential.

6. Conclusions

In this brief paper we have presented initial results for nonlinear AIMD protocols operating in stochastic environments. Future work will involve extending our results to more realistic network scenarios.

Acknowledgement

This work was supported by SFI PI Award 07/IN.1/1901.

Appendix

Proof of Lemma 2.1. For $k \geq k^*$, let $\mathcal{D}(k)$ denote the set of users that have been dropped at least once during the events k^*, \dots, k . We will show by induction that for all $k > k^*$ and $j \in \mathcal{D}(k-1)$, the quantities $w(k)$ and $t_j(k)$ are independent of $t(k^*)$.

So suppose that this statement holds for some event $k > k^*$. If $i \notin \mathcal{D}(k-1)$ then, user i has not been dropped during events $k^*, \dots, k-1$ and (6) implies that $t_i(k) \geq t_i(k^*) \geq T_{m_i}$. It follows from the fact that a_i is affine after T_{m_i} (recall (7)) that

$$a_i(t_i(k) + T(k)) - a_i(t_i(k)) = \alpha_{H_i} T(k). \quad (40)$$

Hence Eqs. (2)–(4) are independent of $t_i(k)$. Thus, $T(k)$ and $w(k+1)$ are independent of $t_i(k)$ and depend only on $w(k)$ and $t_j(k)$ where $j \in \mathcal{D}(k-1)$. Since, by assumption, $w(k)$ and $t_j(k)$ do not depend on $t(k^*)$ for all $j \in \mathcal{D}(k-1)$ it follows that $T(k)$ and $w(k+1)$ are independent of $t(k^*)$. Consider now any user $j \in \mathcal{D}(k)$. It follows from (5) and (6) that $t_j(k+1)$ depends only on $w(k)$ and $t_j(k)$ where $j \in \mathcal{D}(k-1)$; hence $t_j(k+1)$ is independent of $t(k^*)$.

We now show that the statement holds for $k = k^* + 1$. If user i is not dropped at k^* , $t_i(k^*) \geq T_{m_i}$ and $a_i(t_i(k^*) + T(k^*)) - a_i(t_i(k^*)) = \alpha_{H_i} T(k^*)$. Eqs. (2)–(4) imply that $T(k^*)$ and $w(k^* + 1)$ are independent of $t(k^*)$. If $j \in \mathcal{D}(k^*)$ then (5) now implies that $t_j(k^* + 1)$ is independent of $t(k^*)$. \square

Proof of Lemma 4.2. Suppose user one is dropped at an event k . Then

$$w_1(k+1) = \beta_1 w_1 + a_1(T) \quad (41)$$

$$w_2(k+1) = b_2 w_2 + a_2(d_2 t_2 + T) - a_2(d_2 t_2) \quad (42)$$

$$t_1(k+1) = T \quad (43)$$

$$t_2(k+1) = d_2 t_2 + T. \quad (44)$$

Here $(b_2, d_2) = (\beta_2, 0)$ if user two is dropped while $(b_2, d_2) = (1, 1)$ if user two is not dropped. Considering $\tau = \max\{T_{m_2}, d_2 t_2\}$ and using the convexity of a yield

$$a_2(d_2 t_2 + T) - a_2(d_2 t_2) \leq a_2(\tau + T) - a_2(\tau) = \alpha_{H_2} T. \quad (45)$$

Since $b_2 \leq 1$, relationships (42) and (45) result in

$$w_2(k+1) \leq w_2 + \alpha_{H_2} T. \quad (46)$$

Recalling that $w_1(k+1) + w_2(k+1) = w_1 + w_2 = C$, relationships (41) and (46) imply that

$$a_1(T) + \alpha_{H_2} T \geq (1 - \beta_1) w_1 \geq (1 - \beta_1) w_1 = a_1(T_m) + \alpha_{H_2} T_m,$$

that is, $a_1(T) + \alpha_{H_2} T \geq a_1(T_m) + \alpha_{H_2} T_m$. Since a_1 is increasing and $\alpha_{H_2} > 0$, we must have $T \geq T_m$. Hence $t_i(k+1) \geq T_m$ for $i = 1, 2$. If user two is dropped, the proof proceeds in the same fashion. \square

Proof of Lemma 4.3. Suppose user two is not dropped. Then user one must be dropped; hence

$$w_1(k+1) = \beta_1 w_1 + a_1(T) \quad (47)$$

$$w_2(k+1) = w_2 + a_2(t_2 + T) - a_2(t_2). \quad (48)$$

Consider first the situation in which $w_1 \leq \underline{w}_1$. Since $w_1 + w_2 = C \geq \underline{w}_1 + \underline{w}_2$, we must have $w_2 \geq \underline{w}_2$. Clearly $w_2(k+1) \geq w_2$; hence $w_2(k+1) \geq \underline{w}_2$.

Consider now the situation in which $w_1 \geq \underline{w}_1$. Since user one is dropped, it follows from Lemma 4.2 that $T \geq T_m$; hence (47) implies that

$$w_1(k+1) = \beta_1 w_1 + a_1(T_m) + \alpha_{H_1}(T - T_m). \quad (49)$$

The convexity of a and $t_2 \geq 0$ implies that $a_2(t_2 + T) - a_2(t_2) \geq a_2(T) - a_2(0) = a_2(T) = \alpha_{H_2}(T - T_m) + a_2(T_m)$; hence (48) implies that

$$w_2(k+1) \geq w_2 + a_2(T_m) + \alpha_{H_2}(T - T_m). \quad (50)$$

Combining (49) and (50) to eliminate T and using the identities $w_1(k+1) + w_2(k+1) = w_1 + w_2 = C$ result in

$$\begin{aligned} (\alpha_{H_1} + \alpha_{H_2})w_2(k+1) &\geq (\alpha_{H_1} + \alpha_{H_2})C \\ -(\alpha_{H_1} + \beta_1\alpha_{H_2})w_1 + \alpha_{H_1}a_2(T_m) - \alpha_{H_2}a_1(T_m). \end{aligned}$$

Now use the inequalities $w_1 \leq C$ and $C \geq C_2$, where C_2 is given by (20), to obtain that $w_2(k+1) \geq \underline{w}_2$.

The proof proceeds in the same fashion if user one is not dropped. \square

Proof of Lemma 4.5. Consider the sequence $\mathcal{Q} = \{q_{k^*}, \dots, q_{k^f-1}\}$ in $\{1, 2, 3\}$. Using the following algorithm, we obtain a new sequence $\mathcal{P} = \{p_0, p_1, \dots, p_{l-1}\}$ in $\{1, 2, 3, 4, 5\}$ by replacing selected subsequences of the form $\{1, 2\}$ or $\{2, 1\}$ in \mathcal{Q} with 4 and 5, respectively. The new sequence \mathcal{P} does not contain the subsequences $\{1, 2\}$, $\{2, 1\}$, $\{1, 5\}$ or $\{2, 4\}$. This algorithm is illustrated in (51).

Algorithm. Starting with index $K = k^*$, find the first $k \geq K$ with $\{q_k, q_{k+1}\}$ equal to $\{1, 2\}$ or $\{2, 1\}$. If $\{q_k, q_{k+1}\} = \{1, 2\}$, replace it with 4; otherwise replace it with 5. Set the index K to $k+2$.

$$\begin{array}{ll} \mathcal{Q} & 11322221121 \quad 2232113 \\ \mathcal{P} & 113222541 \quad 223513 \end{array} \quad (51)$$

From the above algorithm, it should be clear that there is a subsequence $\{k_0, k_1, \dots, k_l\}$ of $\{k^*, k^* + 1, \dots, k^f\}$ such that $k^* = k_0, k^f = k_l$ and

$$X(k_{j+1}) = \Psi_{p_j}(X(k_j)) \quad (52)$$

for $j = 0, \dots, l-1$. We now show by induction that

$$X(k_j) \in \mathcal{M}_{p_j} \quad \text{for } j = 0, \dots, l-1. \quad (53)$$

So suppose that $X(k_j) \in \mathcal{M}_{p_j}$ for some j . If $p_j = 3, 4$ or 5 , Lemma 4.4 tells us that $X(k_{j+1}) = \Psi_{p_j}(X(k_j))$ belongs to \mathcal{M} ; hence it is in $\mathcal{M}_{p_{j+1}}$. If $p_j = 1$ then, $X(k_j) \in \mathcal{M}_1$ and Lemma 4.4 tells us that $X(k_{j+1}) = \Psi_1(X(k_j))$ belongs to \mathcal{M}_1 . Since \mathcal{P} does not contain the subsequences $\{1, 2\}$ and $\{1, 5\}$, we must have $p_{j+1} = 1, 3$ or 4 , and therefore $X(k_{j+1}) \in \mathcal{M}_{p_{j+1}}$. A similar reasoning applies if $p_j = 2$. Thus, $X(k_j) \in \mathcal{M}_{p_j}$ implies that $X(k_{j+1}) \in \mathcal{M}_{p_{j+1}}$. Since $X(k_0) = X(k^*) \in \mathcal{M} \subset \mathcal{M}_{p_0}$, we obtain (53) by induction.

It now follows from (52), (53) and (30) that, for $j = 0, \dots, l-1$,

$$PX(k_{j+1}) = P\Psi_{p_j}(X(k_j)) = \Phi_{p_j}(PX(k_j)),$$

that is,

$$w(k_{j+1}) = \Phi_{p_j}(w(k_j)). \quad \square$$

Proof of Lemma 4.6. Part (i). Consider $q = 1$ or 3 . Since user one is dropped in either case, the map $\Psi_q : X \mapsto X'$ is given by

$$\begin{aligned} w'_1 &= \beta_1 w_1 + a_1(T), & t'_1 &= T \\ w'_2 &= b_2 w_2 + a_2(d_2 t_2 + T) - a_2(d_2 t_2), & t'_2 &= d_2 t_2 + T \end{aligned} \quad (54)$$

where T , the inter-congestion time associated with X , is given by $w'_1 + w'_2 = C$. Here $(b_2, d_2) = (1, 1)$ if $q = 1$ while $(b_2, d_2) = (\beta_2, 0)$ if $q = 3$.

Consider any $X, Y \in \mathcal{S}^{\text{ext}}$. With $(w, t) = X$ and $(v, s) = Y$, let $\eta = w - v$ and $\tau = t - s$. Since $\eta_1 + \eta_2 = 0$, we have $\|\eta\| = 2|\eta_1|$; hence

$$\|X - Y\| = \|\eta\| + \|\tau\| = 2|\eta_1| + \|\tau\|. \quad (55)$$

Define $(\eta', \tau') = \Psi_q(X) - \Psi_q(Y)$. Then, $\eta'_1 + \eta'_2 = 0$ implies that

$$\|\Phi_q(X) - \Phi_q(Y)\| = 2|\eta'_1| + \|\tau'\|. \quad (56)$$

Let $\delta T = T - S$ where T and S are the inter-congestion times associated with X and Y , respectively. As a consequence of the properties of a_1 and a_2 , there exist scalars $\alpha_1, \alpha_2, \alpha_3$ such that

$$\begin{aligned} a_1(T) - a_1(S) &= \alpha_1 \delta T \\ a_2(d_2 t_2 + T) - a_2(d_2 s_2 + S) &= \alpha_2(d_2 \tau_2 + \delta T) \end{aligned} \quad (57)$$

$$a_2(d_2 t_2) - a_2(d_2 s_2) = \alpha_3 d_2 \tau_2$$

with

$$\alpha_{L_1} \leq \alpha_1 \leq \alpha_{H_1} \quad \text{and} \quad \alpha_{L_2} \leq \alpha_2, \alpha_3 \leq \alpha_{H_2}.$$

To see this consider, for example,

$$\alpha_1 = \begin{cases} \frac{a_1(T) - a_1(S)}{T - S} & \text{if } T \neq S \\ \alpha_{H_1} & \text{if } T = S. \end{cases} \quad (58)$$

The convexity of a_2 implies that $\alpha_3 \leq \alpha_2$. It now follows from description (54) of Ψ_q that

$$\begin{aligned} \eta'_1 &= \beta_1 \eta_1 + \alpha_1 \delta T, & \tau'_1 &= \delta T \\ \eta'_2 &= b_2 \eta_2 + \alpha_2 \delta T + \gamma d_2 \tau_2, & \tau'_2 &= d_2 \tau_2 + \delta T \end{aligned} \quad (59)$$

where $\gamma = \alpha_2 - \alpha_3 \geq 0$. Since $\eta'_1 + \eta'_2 = \eta_1 + \eta_2 = 0$, δT is given by

$$(\beta_1 - b_2)\eta_1 + (\alpha_1 + \alpha_2)\delta T + \gamma d_2 \tau_2 = 0.$$

Solving for δT and substituting into (59) yield

$$\eta'_1 = \left(\frac{\alpha_2 \beta_1 + \alpha_1 b_2}{\alpha_1 + \alpha_2} \right) \eta_1 - d_2 \left(\frac{\gamma \alpha_1}{\alpha_1 + \alpha_2} \right) \tau_2 \quad (60)$$

$$\tau'_1 = \left(\frac{b_2 - \beta_1}{\alpha_1 + \alpha_2} \right) \eta_1 \quad (61)$$

$$\tau'_2 = \left(\frac{b_2 - \beta_1}{\alpha_1 + \alpha_2} \right) \eta_1 + d_2 \left(1 - \frac{\gamma}{\alpha_1 + \alpha_2} \right) \tau_2. \quad (62)$$

As a consequence of the upper and lower bounds on $\alpha_1, \alpha_2, \alpha_3$ there exists a constant c_{13} independent of X and Y such that

$$2|\eta'_1| + \|\tau'\| \leq c_{13}(2|\eta_1| + \|\tau\|). \quad (63)$$

Hence, $\|\Psi_q(X) - \Psi_q(Y)\| \leq c_{13}\|X - Y\|$. One can follow the same procedure to obtain a corresponding constant c_{23} for $q = 2, 3$. Finally, let $c_1 = \max\{c_{13}, c_{23}\}$ to obtain the desired result.

Part (ii). Consider first $q = 1$ or 3 and recall that $\Phi_q(w) = P\Psi_q(w, T_m, T_m)$. Hence $\|\Phi_q(w) - \Phi_q(v)\| = \|\eta'\| = 2|\eta'_1|$. When $q = 3$ we have $d_2 = 0$. When $q = 1$, $d_2 = 1$. Recall that a_2 is affine with slope α_{H_2} after $T_{m_2} \leq T_m$. Hence, when $q = 1$ and $t_2 = s_2 = T_m$, Eq. (57) tells us that $\alpha_2 = \alpha_3 = \alpha_{H_2}$; hence $\gamma = 0$. In either case, it follows from (60) that

$$\eta'_1 = \left(\frac{\alpha_2 \beta_1 + \alpha_1 b_2}{\alpha_1 + \alpha_2} \right) \eta_1. \quad (64)$$

Since $\beta_2 \leq 1$, $\beta_1 < 1$, $0 < \alpha_1 \leq \alpha_{H_1}$ and $0 < \alpha_{L_2} \leq \alpha_2$, we must have

$$\frac{\alpha_2 \beta_1 + \alpha_1 \beta_2}{\alpha_1 + \alpha_2} \leq \frac{\alpha_2 \beta_1 + \alpha_1}{\alpha_1 + \alpha_2} \leq \frac{\alpha_{L_2} \beta_1 + \alpha_{H_1}}{\alpha_{H_1} + \alpha_{L_2}} \leq r$$

and it follows from (64) that $|\eta'_1| \leq r|\eta_1|$. Since $\|\Phi_q(w) - \Phi_q(v)\| = 2|\eta'_1|$ and $\|w - v\| = 2|\eta_1|$, we obtain the desired result that $\|\Phi_q(w) - \Phi_q(v)\| \leq r\|w - v\|$. The proof for $q = 2$ or 3 is similar. \square

Proof of Lemma 4.7. The map $\Phi_4 : w \mapsto w''$ is given by

$$\begin{aligned} w'_1 &= \beta_1 w_1 + a_1(T), & w''_1 &= w'_1 + a_1(T' + T) - a_1(T) \\ w'_2 &= w_2 + \alpha_{H_2} T, & w''_2 &= \beta_2 w'_2 + a_2(T') \end{aligned} \quad (65)$$

where the inter-congestion times T and T' are calculated by imposing the conditions $w'_1 + w'_2 = w''_1 + w''_2 = C$. Also, Corollary 4.2 tells us that $T' \geq T_m$. For any $w, v \in \mathcal{S}$, let $\eta = w - v$ and $\eta'' = \Phi_4(w) - \Phi_4(v)$. The conditions $\eta_1 + \eta_2 = 0$ and $\eta''_1 + \eta''_2 = 0$ yield

$$\|w - v\| = 2|\eta_1| \quad \text{and} \quad \|\Phi_4(w) - \Phi_4(v)\| = 2|\eta''_1|. \quad (66)$$

Let α_1 be given by (58) where T, T' and S, S' are the inter-congestion times associated with w and v , respectively. Convexity of $a(\cdot)$ implies that $\alpha_1 \leq \alpha_H$. The description of Φ_4 in (65) now yields

$$\eta'_1 = \beta_1 \eta_1 + \alpha_1 \delta T \quad (67)$$

$$\eta'_2 = \eta_2 + \alpha_{H_2} \delta T \quad (68)$$

$$\eta''_1 = \eta'_1 + \alpha_{H_1} (\delta T' + \delta T) - \alpha_1 \delta T \quad (69)$$

$$\eta''_2 = \beta_2 \eta'_2 + \alpha_{H_2} \delta T' \quad (70)$$

where $\delta T := T - S$ and $\delta T' := T' - S'$. Since $\eta_1 + \eta_2 = \eta'_1 + \eta'_2 = 0$, Eqs. (67) and (68) imply that

$$\delta T = \left(\frac{1 - \beta_1}{\alpha_1 + \alpha_{H_2}} \right) \eta_1, \quad \eta'_1 = \left(\frac{\alpha_1 + \alpha_{H_2} \beta_1}{\alpha_1 + \alpha_{H_2}} \right) \eta_1. \quad (71)$$

Using $\eta'_1 + \eta'_2 = \eta''_1 + \eta''_2 = 0$, Eqs. (69) and (70) result in

$$\eta''_1 = \left(\frac{\alpha_{H_2} + \alpha_{H_1} \beta_2}{\alpha_{H_1} + \alpha_{H_2}} \right) \eta'_1 + \left(\frac{\alpha_{H_2} (\alpha_{H_1} - \alpha_1)}{\alpha_{H_1} + \alpha_{H_2}} \right) \delta T.$$

Substituting the previously obtained expressions (71) for η'_1 and δT results in $\eta''_1 = \rho \eta_1$ where

$$\rho = \frac{\alpha_1 (\alpha_{H_1} \beta_2 + \alpha_{H_2} \beta_1) + \alpha_{H_2} (\alpha_{H_1} (1 + \beta_1 \beta_2 - \beta_1) + \alpha_{H_2} \beta_1)}{(\alpha_1 + \alpha_{H_2}) (\alpha_{H_1} + \alpha_{H_2})}.$$

Since $1 + \beta_1 \beta_2 - \beta_1 \geq \beta_2$, the above expression for ρ achieves its maximum value when α_1 is at its minimum value of α_{L_1} ; the corresponding maximum value of ρ is r_4 . Hence $|\eta''_1| \leq r_4 |\eta_1| \leq r |\eta_1|$. Thus $\|\Phi_4(w) - \Phi_4(v)\| \leq r \|w - v\|$. The map Φ_5 is handled similarly. \square

References

Baccelli, F., & Hong, D. (2002). AIMD, fairness and fractal scaling of TCP traffic. In *Proc. of IEEE INFOCOM 2002* (pp. 229–238).

Barnsley, M., Demko, S., Elton, J., & Geronimo, J. (1988). Invariant measures for markov processes arising from iterated function systems with place dependent probabilities. *Annales de l'Institut Henri Poincaré Probabilités et Statistiques*, 24, 367–394.

Chiu, D., & Jain, R. (1989). Analysis of the increase/decrease algorithms for congestion avoidance in computer networks. *Journal of Computer Networks*, 17, 1–14.

Floyd, S. (2003). High speed TCP for large congestion windows. *Tech. rep. Internet draft draft-floyd-tcp-highspeed-02.txt*.

Jacobson, V. (1988). Congestion avoidance and control. In *Proc. of SIGCOMM*.

Jin, C., Wei, D., & Low, S. (2003). FAST TCP: Motivation, architecture, algorithms, performance. *Caltech CS Report CaltechCSTR:2003:010*.

Kelly, T. (2002). On engineering a stable and scalable TCP variant. *Tech. rep. Cambridge University Engineering Department Technical Report CUED/F-INFENG/TR.435*.

King, C., Shorten, R., Wirth, F., & Akar, M. (2008). Growth conditions for the global stability of high-speed communication networks with a single congested link. *IEEE Transactions on Automatic Control*, 53(7), 1770–1774.

Low, S., Paganini, F., & Doyle, J. (2002). Internet congestion control. *IEEE Control Systems Magazine*, 32(1), 28–43.

Rothblum, U., & Shorten, R. (2007). Nonlinear AIMD congestion control and contraction mappings. *SIAM Journal of Control and Optimisation*, 46(5), 1882–1896.

Shorten, R., King, C., Wirth, F., & Leith, D. (2007). Modelling TCP in droptail environments. *Automatica*, 43, 441–449.

Shorten, R., Wirth, F., & Leith, D. (2006). A positive systems model of TCP-like congestion control: Asymptotic analysis. *IEEE/ACM Transactions on Networking*, 14, 616–629.

Stenflo, O. (2002). Uniqueness of invariant measures for place dependent iterations of functions. *IMA Journal of Applied Mathematics*, 132, 13–32.

Xu, L., Harfoush, K., & Rhee, I. (2004). Binary increase congestion control (BIC) for fast long-distance networks. In *Proc. of IEEE INFOCOM 2004* (pp. 2514–2524).



Martin Corless is currently a Professor in the School of Aeronautics & Astronautics at Purdue University, West Lafayette, Indiana, USA. He received a B.E. from University College Dublin, Ireland and a Ph.D. from the University of California at Berkeley; both degrees are in mechanical engineering. He is the recipient of a National Science Foundation Presidential Young Investigator Award. His research is concerned with obtaining tools which are useful in the robust analysis and control of systems containing significant uncertainty and in applying these results to aerospace, mechanical and communication systems.



Chris King obtained his PhD in Theoretical Physics at Harvard University in 1984. He has been with the Department of Mathematics at Northeastern University since 1991. His research interests include mathematical physics, Quantum Information Theory, control theory, and network analysis.



Robert Shorten graduated from the University College Dublin (UCD) in 1990 with a First Class Honours B.E. degree in Electronic Engineering. He was awarded a M.Eng.Sc in 1993, and a Ph.D. in 1996, both from UCD. From 1993 to 1996 he was the holder of a Marie Curie Fellowship to conduct research at the Daimler-Benz Research Institute for Information Technology in Berlin. In 1996 he was invited as a visiting fellow to the Center for Systems Science, Yale University, to work with Professor K.S. Narendra. Since returning to Ireland in 1997 as the recipient of a European Presidential Fellowship, he has built up a research group in the area of switching systems. He is a member of the Editorial Board for the IET Journal of Control Theory & Applications and is a co-founder of the Hamilton Institute, NUI Maynooth.